

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD19 Monoclonal

Antibody(PE/Cyanine7 Conjugated)

Catalog: DL20019F

Synonyms: B-lymphocyte antigen CD19,Cd19,Differentiation

antigen CD19,CD19

Background: CD19 is a 95 kD glycoprotein also known as B4. It is

> a member of the Ig superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81,

forms a molecular complex integral to B cell

activation.

Form: Liquid **Isotype:** Rat IgG2a, κ



Size: 25µg/100µg Host: Rat

Reactivity: Mouse **Application:** FCM

Concentration: 0.2 mg/mL Conjugation: PE/Cyanine7

SwissProt: P25918

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in 100 μL volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD19 Monoclonal Antibody(PE/Cyanine7 Conjugated)[Used at 0.05 μg/10⁶ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.





Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.