

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-8	2732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD19 Monoclonal Antibody(AF488		
	Conjugated)		
Catalog:	DL20022F		
Synonyms:	B-lymphocyte antigen CD19,Cd19,Differentiation antigen CD19,CD19		
Background:	CD19 is a 95 kD glycoprotein also known as B4. It is a member of the Ig superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.		
Form: Liquid	<b>Isotype:</b> Rat IgG2a, κ		



Size: 25µg/100µg		Host: Rat	
Reactivity: Mouse		Application: FCM	
Concentration: 0.5 mg/mL		Conjugation: AF488	
SwissProt:	P25918		
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		


## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup>cells in 100  $\mu$ L volume].

## Data:

Mouse splenocytes are stained with Anti-Mouse CD19 Monoclonal Antibody(AF488 Conjugated)[Used at  $0.2 \mu g/10^6$  cells dilution](filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.

## **DEVELOP**



## Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

