

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn) [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Anti-Mouse CD161/NK1.1 Monoclonal  
Antibody(AF488 Conjugated)

**Catalog:** DL20035F

**Synonyms:** Killer cell lectin-like receptor subfamily B member  
1C, Klrb1c, CD161 antigen-like family member  
C, Ly-55c, CD161/NK1.1, NKR-P1.9, NKR-P1C, NKR  
-P1 40, CD161c

**Background:** NK-1.1 surface antigen, also known as  
CD161b/CD161c and Ly-55, is encoded by the  
NKR- P1B/NKR-P1C gene. It is expressed on NK  
cells and NK-T cells in some mouse strains,  
including C57BL/6, FVB/N, and NZB, but not AKR,  
BALB/c, CBA/J, C3H, DBA/1, DBA/2, NOD, SJL,  
and 129. Expression of NKR-P1C antigen has been  
correlated with lysis of tumor cells in vitro and

rejection of bone marrow allografts in vivo. NK-1.1 has also been shown to play a role in NK cell activation, IFN- $\gamma$  production, and cytotoxic granule release. NK-1.1 and DX5 are commonly used as mouse NK cell markers.

**Form:** Liquid

**Isotype:** Mouse IgG2a,  $\kappa$

**Size:**

50Tests/100Tests/100Tests $\times$ 2

**Host:** Mouse

**Reactivity:** Mouse

**Application:** FCM

**Concentration:** 5  $\mu$ L

**Conjugation:** AF488

**SwissProt:** P27814

**Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.


### Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined

for individual use.

### Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD161/NK1.1 Monoclonal Antibody(AF488 Conjugated) and Anti-Mouse CD49b Monoclonal Antibody(APC Conjugated)(Left). Unstained splenocytes are used as control(Right).



### Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.