

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

**Product name:** Anti-Mouse CD161/NK1.1 Monoclonal

Antibody(AF647 Conjugated)

Catalog: DL20036F

**Synonyms:** Killer cell lectin-like receptor subfamily B member

1C,Klrb1c,CD161 antigen-like family member

C,Ly-55c,CD161/NK1.1,NKR-P1.9,NKR-P1C,NKR

-P1 40,CD161c

**Background:** NK-1.1 surface antigen, also known as

CD161b/CD161c and Ly-55, is encoded by the

NKR-P1B/NKR-P1C gene. It is expressed on NK

cells and NK-T cells in some mouse strains,

including C57BL/6, FVB/N, and NZB, but not AKR, BALB/c, CBA/J, C3H, DBA/1, DBA/2, NOD, SJL, and 129. Expression of NKR-P1C antigen has been correlated with lysis of tumor cells in vitro and



rejection of bone marrow allografts in vivo. NK-1.1 has also been shown to play a role in NK cell activation, IFN-y production, and cytotoxic granule release. NK-1.1 and DX5 are commonly used as mouse NK cell markers.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: Host: Mouse

50Tests/100Tests/100Tests×2

**Reactivity:** Mouse **Application:** FCM

Concentration: 5 µL **Conjugation:** AF647

SwissProt: P27814

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

## Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined



for individual use.

## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD161/NK1.1 Monoclonal Antibody(AF647 Conjugated) and Anti-Mouse CD49b Monoclonal Antibody(AF488 Conjugated)(left). Splenocytes stained with Anti-Mouse CD49b Monoclonal Antibody(AF488 Conjugated)(right) are used as control.



## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.