

Manufacturer/Supplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China		
TELE: 86-510-82732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>		
Email: info@dldevelop.com.cn service@dldevelop.com.cn		
Product name:	Anti-Mouse MHC II (I-A/I-E) Monoclonal	
	Antibody(PE Conjugated)	
Catalog:	DL20051F	
Synonyms:	H2-Ab1/Eb1, Major histocompatibility protein class	
e e	II beta chain,MHC class II H2-IA-beta-psi,I-E beta	
	MHC class II,MHC class II	
Background:	These class II molecules are expressed on antigen	
	presenting cells (including B cells) and a subset of T	
	cells from H-2b,d,q,r bearing mice and are involved	
	in antigen presentation to T cells expressing	
	CD3/TCR and CD4 proteins.	
	ebs/rect and eb+ proteins.	
Form: Liquid	<b>Isotype:</b> Rat IgG2b, κ	
Torme Diquid	1900 per 14t 15020, K	
Size:	Host: Rat	



50Tests/100Tests/100Tests×2 Reactivity: Mouse

**Application:** FCM

**Concentration:** 5 µL

**Conjugation: PE** 

SwissProt:	P14483,O78196
Buffer:	Phosphate buffered solution, pH 7.2, containing
	0.09% stabilizer and 1% protein protectant.


## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Data:

Mouse splenocytes are stained with Anti-Mouse I-A/I-E Monoclonal Antibody(PE Conjugated)(filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.

## **DEVELOP**



## Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

