

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: AF/LE Purified Anti-Mouse CD86 Monoclonal
Antibody

Catalog: DL20099F

Synonyms: T-lymphocyte activation antigen
CD86, Cd86, Activation B7-2 antigen, Early T-cell
costimulatory molecule 1, ETC-1

Background: CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is a ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86

binds to CD28 to transduce co- stimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

Form: Liquid

Isotype: Rat IgG2a, κ

Size: 50 μ g/500 μ g/1mg

Host: Rat

Reactivity: Mouse

Application: FCM,IHC-F,IP,ICC

Concentration: 0.5 mg/mL

Conjugation: None (Purified antibody,Azide Free/Low endotoxin)

SwissProt: P42082

Buffer: 0.2 μ m filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu$ g per 10^6 cells in 100 μ L volume or 100 μ L of

whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:

Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.