

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD86 Monoclonal Antibody(Biotin

Conjugated)

Catalog: DL20101F

Synonyms: T-lymphocyte activation antigen

CD86,Cd86,Activation B7-2 antigen,Early T-cell

costimulatory molecule 1,ETC-1

Background: CD86 is an 80 kD immunoglobulin superfamily

member also known as B7-2, B70, and Ly-58. CD86

is expressed on activated B and T cells,

macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is a ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class- switching and triggering of NK cell-mediated cytotoxicity. CD86



binds to CD28 to transduce co-stimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

Form: Liquid Isotype: Rat IgG2a, κ

Size: $25\mu g/100\mu g$ Host: Rat

Reactivity: Mouse **Application:** FCM

Concentration: 0.5 mg/mL **Conjugation:** Biotin

SwissProt: P42082

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.