

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse F4/80 Monoclonal Antibody(FITC

Conjugated)

**Catalog:** DL20123F

**Synonyms:** Adhesion G protein-coupled receptor E1, Adgre1, Cell

surface glycoprotein F4/80,EGF-like module

receptor 1, Adgre 1, Emr 1, Gpf 480

**Background:** F4/80 is a 160 kD glycoprotein. It is characterized as

a member of the epidermal growth factor

(EGF)-transmembrane 7 (TM7) family. F4/80, also known as EMR1 or Ly71, has been widely used as a murine macrophage marker, which is expressed on

the majority of tissue macrophages including

peritoneal macrophages, macrophages in lung, gut,

thymus and red pulp of spleen (but not on the macrophages located in T cell areas of the spleen,



lymph node and Peyer's patch), Kuffer cells, Langerhans cells, and bone marrow stromal cells. F4/80 has also been shown on a subset of dendritic cells. The biological ligand of F4/80 has not been identified, but it has been reported that F4/80 is required for induction of CD8+ T cells-mediated peripheral tolerance.

Form: Liquid **Isotype:** Rat IgG2b, κ

Size: Host: Rat

50Tests/100Tests/100Tests×2

**Reactivity:** Mouse **Application:** FCM

Concentration: 5 µL **Conjugation: FITC** 

O61549 SwissProt:

Phosphate buffered solution, pH 7.2, containing **Buffer:** 

0.09% stabilizer and 1% protein protectant.

## Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per



100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Data:

Mouse abdominal macrophages elicited by starch broth are stained with Anti-Mouse F4/80 Monoclonal Antibody(FITC Conjugated)(filled gray histogram). Unstained macrophages (blank black histogram) are used as control.



## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.