

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Anti-Mouse F4/80 Monoclonal Antibody(AF647			
	Conjugated)			
Catalog:	DL20132F			
Synonyms:	Adhesion G protein-coupled receptor E1,Adgre1,Cell surface glycoprotein F4/80,EGF-like module			
	receptor 1,Adgre1,Emr1, Gpf480			
Background: F4/80 is a 160 kD glycoprotein. It is characterized				
a member of the epidermal growth factor (EGF)-transmembrane 7 (TM7) family. F4/80, also				
		known as EMR1 or Ly71, has been widely used as a		
murine macrophage marker, which is expressed on the majority of tissue macrophages including				
		peritoneal macrophages, macrophages in lung, gut,		
	thymus and red pulp of spleen (but not on the			
	macrophages located in T cell areas of the spleen,			



lymph node and Peyer's patch), Kuffer cells,		
Langerhans cells, and bone marrow stromal cells.		
F4/80 has also been shown on a subset of dendritic		
cells. The biological ligand of F4/80 has not been		
identified, but it has been reported that F4/80 is		
required for induction of CD8+ T cells-mediated		
peripheral tolerance.		

Form: Liquid		Isotype: Rat IgG2b, κ		
Size:		Host: Rat		
50Tests/100Te	sts/100Tests×2			
Reactivity: Mouse		Application: FCM		
Concentration: 5 µL		Conjugation: AF647		
SwissProt:	Q61549			
Buffer:	-	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per



 $100 \ \mu L$ of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Mouse abdominal macrophages elicited by starch broth are stained with Anti-Mouse F4/80 Monoclonal Antibody(AF647 Conjugated)(filled gray histogram). Unstained macrophages (blank black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

