

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD16/32 Monoclonal Antibody(PerCP Conjugated)

Catalog: DL20149F

Synonyms: CD16a/b, CD32, CD32A/B, Fc fragment of IgG low affinity IIa/b receptor, Fc fragment of IgG low affinity IIIa/b receptor, Fc fragment of IgG low affinity IIIb receptor, Fc gamma receptor III A/B, FCGR2A, FcGR, FCGR2A/BFCGR3, FCGR3A/B, Fc gamma RIIa/b

Background: CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes

and mediate adaptive immune responses.

Form: Liquid

Isotype: Rat IgG2b, κ

Size: 20 Tests/50 Tests/100 Tests/200 Tests

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration:

Conjugation: PerCP

SwissProt: P08101,P08508,Q5D5I8,Q5D5J5,Q7TMW9,Q9ES9
2

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:



Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.