

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-801				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Anti-Mouse CD16/32 Monoclonal Antibody(APC			
	Conjugated)			
Catalog:	DL20158F			
0				
Synonyms:	CD16a/b,CD32,CD32A/B,Fc fragment of IgG low affinity IIa/b receptor,Fc fragment of IgG low			
<i>. .</i>				
	affinity IIIa/b receptor,Fc fragment of IgG low			
	affinity IIIb receptor,Fc gamma receptor III			
	A/B,FCG2A,FcGR,FCGR2A/BFCGR3,FCGR3A/B,			
	Fc gamma RIIa/b			
	r o guillina rera o			
Background:	CD16 is low affinity IgG Fc receptor III (FcR III)			
Dackground.	and CD32 is FcR II. CD16/CD32 are expressed on B			
	-			
cells, monocytes/macrophages, NK cells,				
	granulocytes, mast cells, and dendritic cells. The Fc			
	receptors bind antibody-antigen immune complexes			

www.dldevelop.com



and mediate adaptive immune responses.

Form: Liquid		Isotype: Rat IgG2b, κ
Size: 25µg/100µg		Host: Rat
Reactivity: Mouse		Application: FCM
Concentration: 0.2 mg/mL		Conjugation: APC
SwissProt:	P08101,P08508,C	Q5D518,Q5D5J5,Q7TMW9,Q9ES9
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	

## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup>cells in 100  $\mu$ L volume].


## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse

www.dldevelop.com



CD16/32 Monoclonal Antibody(APC Conjugated)[Used at  $0.02 \mu g/10^6$  cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



## Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

