

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD16/32 Monoclonal Antibody(AF488
Conjugated)

Catalog: DL20164F

Synonyms: CD16a/b, CD32, CD32A/B, Fc fragment of IgG low
affinity IIa/b receptor, Fc fragment of IgG low
affinity IIIa/b receptor, Fc fragment of IgG low
affinity IIIb receptor, Fc gamma receptor III
A/B, FCGR2A, FcGR, FCGR2A/BFCGR3, FCGR3A/B,
Fc gamma RIIa/b

Background: CD16 is low affinity IgG Fc receptor III (FcR III)
and CD32 is FcR II. CD16/CD32 are expressed on B
cells, monocytes/macrophages, NK cells,
granulocytes, mast cells, and dendritic cells. The Fc
receptors bind antibody-antigen immune complexes

and mediate adaptive immune responses.

Form: Liquid

Isotype: Rat IgG2b, κ

Size: 25 μ g/100 μ g

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration: 0.5 mg/mL

Conjugation: AF488

SwissProt: P08101,P08508,Q5D5I8,Q5D5J5,Q7TMW9,Q9ES9
2

Buffer: Phosphate buffered solution, pH 7.2, containing
0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶cells in 100 μ L volume].

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.