

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD29 Monoclonal Antibody(FITC
Conjugated)

Catalog: DL20169F

Synonyms: Integrin beta-1, ITB1, Itgb1, Fibronectin receptor
subunit beta, VLA-4 subunit beta, CD29

Background: The KMI6 antibody reacts with the 130-kDa integrin $\beta 1$ chain (CD29). CD29 is expressed on the cell surface as a heterodimer with one of the distinct integrin α chains. With $\alpha 1$ through $\alpha 6$ (CD49a through CD49f), it forms the VLA-1 through VLA-6 complexes, respectively, and with αv (CD51), it forms $\alpha v\beta 1$ integrin. It also associates with the integrin $\alpha 7$, $\alpha 8$, and $\alpha 9$ chains in non-lymphoid tissues. As a result, CD29 has a broad tissue distribution, including leukocytes, endothelia,

smooth muscle, epithelia, and oocytes. The epitope recognized by the KMI6 mAb can be detected by immunoblot in unreduced extracts from a variety of cells and tissues having a number of different β 1-integrin heterodimer combinations. However, the epitope is not well expressed on the surface of the majority of hematopoietic cells. It is detected on the surface of fetal liver cells, some bone marrow stromal cells, and some cultured hematopoietic cell lines, including P815 and PA6. Surface expression of the CD29 epitope recognized by mAb KMI6 on thymocytes and T lymphocytes appears to be developmentally regulated, and KMI6 antibody enhances the adhesion of KMI6 thymocytes to fibronectin and laminin.

Form: Liquid

Isotype: Rat IgG2a, κ

Size: 50 Tests/100 Tests/500 Tests

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration:

Conjugation: FITC

SwissProt: P09055

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

C57BL/6 murine bone marrow cells are stained with Anti-Mouse CD29 Monoclonal Antibody(FITC Conjugated)(filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.