

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD16 Monoclonal Antibody(Biotin
Conjugated)

Catalog: DL20201F

Synonyms: Low affinity immunoglobulin gamma Fc region
receptor III-A, FCGR3A, CD16a antigen, Fc-gamma
RIII-alpha, Fc-gamma RIII, Fc-gamma
RIIIa, FcRIII, FcRIIIa, FcR-10, IgG Fc receptor
III-2, CD16a

Background: CD16 is known as low affinity IgG receptor III
(FcγRIII). It is expressed as two distinct forms
(CD16a and CD16b). CD16a (FcγRIIIA) is a 50-65
kD polypeptide-anchored transmembrane protein. It
is expressed on the surface of NK cells, activated
monocytes, macrophages, and placental trophoblasts
in humans. CD16b (FcγRIIIB) is a 48 kD

glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

Form: Liquid

Isotype: Mouse IgG1, κ

Size: 25 μ g/100 μ g

Host: Mouse

Reactivity: Human

Application: FCM

Concentration: 0.5 mg/mL

Conjugation: Biotin

SwissProt: P08637

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 2.0 μ g per 10^6 cells in 100 μ l volume or 100 μ l of whole blood. It is recommended

that the reagent be titrated for optimal performance for each application.

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.