

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Anti-Human CD16 Monoclonal  
Antibody(PE/Cyanine5 Conjugated)

**Catalog:** DL20206F

**Synonyms:** Low affinity immunoglobulin gamma Fc region  
receptor III-A, FCGR3A, CD16a antigen, Fc-gamma  
RIII-alpha, Fc-gamma RIII, Fc-gamma  
RIIIa, FcRIII, FcRIIIa, FcR-10, IgG Fc receptor  
III-2, CD16a

**Background:** CD16 is known as low affinity IgG receptor III  
(FcγRIII). It is expressed as two distinct forms  
(CD16a and CD16b). CD16a (FcγRIIIA) is a 50-65  
kD polypeptide-anchored transmembrane protein. It  
is expressed on the surface of NK cells, activated  
monocytes, macrophages, and placental trophoblasts  
in humans. CD16b (FcγRIIIB) is a 48 kD

glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:** 20 Tests/50 Tests/100 Tests/200 Tests

**Host:** Mouse

**Reactivity:** Human

**Application:** FCM

**Concentration:**

**Conjugation:** PE/Cyanine5

**SwissProt:** P08637

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

### **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ l staining volume or per 100  $\mu$ l of whole blood). Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Data:**

Human peripheral blood cells are stained with Anti-Human CD16 Monoclonal Antibody(PE/Cyanine5 Conjugated)(filled gray histogram). Unstained peripheral blood cells (blank black histogram) are used as control.

#### **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.