

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD16 Monoclonal Antibody(AF488
Conjugated)

Catalog: DL20210F

Synonyms: Low affinity immunoglobulin gamma Fc region
receptor III-A, FCGR3A, CD16a antigen, Fc-gamma
RIII-alpha, Fc-gamma RIII, Fc-gamma
RIIIa, FcRIII, FcRIIIa, FcR-10, IgG Fc receptor
III-2, CD16a

Background: CD16 is known as low affinity IgG receptor III
(FcγRIII). It is expressed as two distinct forms
(CD16a and CD16b). CD16a (FcγRIIIA) is a 50-65
kD polypeptide-anchored transmembrane protein. It
is expressed on the surface of NK cells, activated
monocytes, macrophages, and placental trophoblasts
in humans. CD16b (FcγRIIIB) is a 48 kD

glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

Form: Liquid

Isotype: Mouse IgG1, κ

Size: 20 Tests/50 Tests/100 Tests/200 Tests

Host: Mouse

Reactivity: Human

Application: FCM

Concentration:

Conjugation: AF488

SwissProt: P08637

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Human peripheral blood cells are stained with Anti-Human CD16 Monoclonal Antibody(AF488 Conjugated)(filled gray histogram). Unstained peripheral blood cells (blank black histogram) are used as control.

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.