

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD56 Monoclonal

Antibody(PE/Cyanine5 Conjugated)

Catalog: DL20216F

Synonyms: Neural cell adhesion molecule

1,NCAM1,N-CAM-1,NCAM-1,NCAM

Background: CD56 is a single transmembrane glycoprotein also

known as NCAM (neural cell adhesion molecule),

Leu-19, or NKH1. It is a member of the Ig

superfamily. The 140 kD isoform is expressed on NK and NKT cells. CD56 is also expressed in the brain

(cerebellum and cortex) and at neuromuscular

junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias

also express CD56. CD56 plays a role in homophilic



and heterophilic adhesion via binding to itself or heparan sulfate.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: 20 Tests/50 Tests/100 Host: Mouse

Tests/200 Tests

Application: FCM **Reactivity:** Human

Concentration: Conjugation: PE/Cyanine5

SwissProt: P13591

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µl staining volume or per 100 µl of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:



Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.