

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD86 Monoclonal Antibody(PerCP

Conjugated)

Catalog: DL20256F

Synonyms: T-lymphocyte activation antigen

CD86,Cd86,Activation B7-2 antigen,Early T-cell

costimulatory molecule 1,ETC-1

Background: CD86 is an 80 kD immunoglobulin superfamily

member also known as B7-2, B70, and Ly-58. CD86

is expressed on activated B and T cells,

monocytes/macrophages, dendritic cells, and

astrocytes. CD86, along with CD80, is the ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated



cytotoxicity. CD86 binds to CD28 to transduce costimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can bind to CD152 as well, also known as CTLA-4, to deliver an inhibitory signal to T cells.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: 20 Tests/50 Tests/100 Host: Mouse

Tests/200 Tests

Reactivity: Human **Application:** FCM

Conjugation: PerCP Concentration:

Swiss Prot P42081

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.