

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: AF/LE Purified Anti-Mouse CD3 Monoclonal
Antibody

Catalog: DL20263F

Synonyms: T-cell surface glycoprotein CD
3epsilon/delta/gamma/zeta
chain, CD3E/D/G/Z, CD3e/d/g/z, CD3E/D/G/Z, CD3

Background: CD3, also known as T3, is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 is composed of CD3ε, δ, γ and ζ chains. It forms a TCR complex by associating with TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.

Form: Liquid

Isotype: Rat IgG2b, κ

Size: 50 μ g/500 μ g/1mg

Host: Rat

Reactivity: Mouse

Application: FCM,IHC-F,IP,ICC

Concentration: 0.5 mg/mL

Conjugation: None (Purified antibody,Azide Free/Low endotoxin)

SwissProt: P04235,P11942,P22646,P24161,P29020

Buffer: 0.2 μ m filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.25 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:



Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.