

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: AF/LE Purified Anti-Mouse CD132 Monoclonal

Antibody

**Catalog:** DL20286F

Cytokine receptor common subunit **Synonyms:** 

gamma, Il 2rg, Interleukin-2 receptor subunit

gamma,IL-2 receptor subunit gamma,IL-2R subunit

gamma, IL-2RG, gammaC, p64, CD132

**Background:** CD132 is a 64-70 kD type I transmembrane

> glycoprotein of the Ig superfamily, and is also known as common γ chain (γc) or IL-2 receptor γ subunit. It is expressed broadly on T- and B- lymphocytes, NK cells, monocytes, and granulocytes. CD132 is an essential component of cytokine receptors for IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21. Ligand binding induces tyrosine phosphorylation and initiates



signaling through a JAK/STAT pathway. CD132 mutation results in X-linked severe combined immune deficiency (XSCID).

Form: Liquid **Isotype:** Rat IgG2b, κ

**Size:** 50µg/500µg/1mg Host: Rat

**Reactivity:** Mouse **Application:** FCM,IP

Concentration: 0.5 mg/mL Conjugation: None (Purified

antibody, Azide Free/Low

endotoxin)

SwissProt: P34902

**Buffer:** 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low

Endotoxin (LE): Contains no stabilizers or

## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \,\mu g$  per  $10^6$  cells in 100  $\mu L$  volume or 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



## **Storage:**

Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.