

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD132 Monoclonal Antibody(APC
Conjugated)

Catalog: DL20292F

Synonyms: Cytokine receptor common subunit
gamma, IL2rg, Interleukin-2 receptor subunit
gamma, IL-2 receptor subunit gamma, IL-2R subunit
gamma, IL-2RG, gammaC, p64, CD132

Background: CD132 is a 64-70 kD type I transmembrane glycoprotein of the Ig superfamily, and is also known as common γ chain (γ_c) or IL-2 receptor γ subunit. It is expressed broadly on T- and B- lymphocytes, NK cells, monocytes, and granulocytes. CD132 is an essential component of cytokine receptors for IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21. Ligand binding induces tyrosine phosphorylation and initiates

signaling through a JAK/STAT pathway. CD132 mutation results in X-linked severe combined immune deficiency (XSCID).

Form: Liquid

Isotype: Rat IgG2b, κ

Size: 25 μ g/50 μ g/100 μ g/200 μ g

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration: 0.2 mg/mL

Conjugation: APC

SwissProt: P34902

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results or refer to the dilution we used in the experiment.

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.