

Manufacturer/Supplier:	
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD	
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District	
Wuxi Jiangsu Province, China	
TELE: 86-510-8	FAX: 86-510-82720101-8014
Web : <u>www.dldevelop.com</u>	
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>	
Product name:	Anti-Mouse CD48 Monoclonal
	Antibody(PE/Cyanine5.5 Conjugated)
Catalog:	DL20325F
Synonyms:	CD48 antigen,Cd48,BCM1 surface
	antigen,BLAST-1,HM48-1,MRC OX-45 surface
	antigen,SLAMF2,sgp-60,CD48
Background:	CD48 is a 45 kD GPI-anchored glycoprotein also
2 weigi o wii w	known as BCM1, Blast-1 (human), and OX-45 (rat).
	It is a member of the Ig superfamily, expressed on T
	and B cells and monocytes/macrophages. It plays a
	role in adhesion and T cell recognition. The primary
	ligands for CD48 are CD2 and CD244.
	-
Form: Liquid	Isotype: Armenian Hamster IgG



Size: 20 Tests/50 Tests/100 Tests/200 Tests Reactivity: Mouse Host: Armenian Hamster

**Application:** FCM

**Concentration:** 

**Conjugation:** PE/Cyanine5.5

SwissProt: P18181

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

## **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.