

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD48 Monoclonal Antibody(FITC
Conjugated)

Catalog: DL20329F

Synonyms: CD48 antigen, Cd48, BCM1 surface
antigen, BLAST-1, HM48-1, MRC OX-45 surface
antigen, SLAMF2, sgp-60, CD48

Background: CD48 is a 45 kD GPI-anchored glycoprotein also
known as BCM1, Blast-1 (human), and OX-45 (rat).
It is a member of the Ig superfamily, expressed on T
and B cells and monocytes/macrophages. It plays a
role in adhesion and T cell recognition. The primary
ligands for CD48 are CD2 and CD244.

Form: Liquid

Isotype: Armenian Hamster IgG

Size: 25µg/100µg

Host: Armenian Hamster

Reactivity: Mouse

Application: FCM

Concentration: 0.5 mg/mL

Conjugation: FITC

SwissProt: P18181


Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10⁶cells in 100 µL volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD48 Monoclonal Antibody(FITC Conjugated)[Used at 0.2 µg/10⁶ cells dilution](filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.