

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Purified Anti-Mouse CD22 Monoclonal Antibody		
Catalog:	DL20363F		
8			
Synonyms:	B-cell receptor CD22,Cd22,B-lymphocyte cell adhesion molecule,BL-CAM,Sialic acid-binding Ig-like lectin 2,Siglec-2,T-cell surface antigen		
	Leu-14,CD22,Lyb-8, Siglec2		
	Leu-14, CD22, Lyb-6, Signe2		
Background:	The Cy34.1 monoclonal antibody specifically binds		
Dackground.			
	to the B-lymphocyte differentiation antigen CD22 on		
	strains having the Lyb-8.2 alloantigen (e.g., A,		
	BALB/c, CBA, C3H/He, C57BL, C57L, C58, SJL,		
	SWR, but not AKR, DBA/1, DBA/2, NZB, PL).		
	CD22 is expressed at high levels on mature		
	peripheral B lymphocytes (follicular and marginal		
	zone), B-1 cells (CD5+ B cells), and plasma cells. It		
	is a member of the Ig gene superfamily and		

DEVELOP

associates with the B-cell antigen receptor. Its sialic acid- binding immunoglobulin-like lectin (siglec) extracellular region mediates B-cell adhesion to ligands on endothelial cells in the bone marrow. Its intracellular domain is phosphorylated after cross-linking of antigen receptor or MHC class II antigen. It is involved in negative regulation of B-cell activation and protection from autoimmunity. B-cell proliferative responses to LPS or anti-mouse Ig μ chain are augmented in the presence of Cy34.1 mAb.

Form: Liquid		Isotype: Mouse IgG1, κ	
Size: 25µg/100µg		Host: Mouse	
Reactivity: Mouse		Application: FCM,IP	
Concentration: 0.5 mg/mL		Conjugation: Unconjugated	
SwissProt:	P35329		
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.		
Recommended Use:			



Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.25 \ \mu g \ per \ 10^6 \ cells$ in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:

Keep as concentrated solution. Store at 2~8°C. Do not freeze. This product is guaranteed up to one year from purchase.

