

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Anti-Mouse CD22 Monoclonal  
Antibody(PE/Cyanine5 Conjugated)

**Catalog:** DL20368F

**Synonyms:** B-cell receptor CD22, Cd22, B-lymphocyte cell  
adhesion molecule, BL-CAM, Sialic acid-binding  
Ig-like lectin 2, Siglec-2, T-cell surface antigen  
Leu-14, CD22, Lyb-8, Siglec2

**Background:** The Cy34.1 monoclonal antibody specifically binds  
to the B-lymphocyte differentiation antigen CD22 on  
strains having the Lyb-8.2 alloantigen (e.g., A,  
BALB/c, CBA, C3H/He, C57BL, C57L, C58, SJL,  
SWR, but not AKR, DBA/1, DBA/2, NZB, PL).  
CD22 is expressed at high levels on mature  
peripheral B lymphocytes (follicular and marginal  
zone), B-1 cells (CD5+ B cells), and plasma cells. It

is a member of the Ig gene superfamily and associates with the B-cell antigen receptor. Its sialic acid-binding immunoglobulin-like lectin (siglec) extracellular region mediates B-cell adhesion to ligands on endothelial cells in the bone marrow. Its intracellular domain is phosphorylated after cross-linking of antigen receptor or MHC class II antigen. It is involved in negative regulation of B-cell activation and protection from autoimmunity. B-cell proliferative responses to LPS or anti-mouse Ig  $\mu$  chain are augmented in the presence of Cy34.1 mAb.

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:**

**Host:** Mouse

50Tests/100Tests/100Tests $\times$ 2

**Reactivity:** Mouse

**Application:** FCM

**Concentration:** 5  $\mu$ L

**Conjugation:** PE/Cyanine5

**SwissProt:** P35329

**Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

### Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.