

Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web: www.dldevelop.com			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD127/IL-7RA Monoclonal		
	Antibody(AF488 Conjugated)		
Catalog:	DL20397F		
Synonyms:	Interleukin-7 receptor subunit alpha,Il7r,IL-7		
	receptor subunit alpha,IL-7R subunit		
	alpha,IL-7R-alpha,IL-7RA,CD127		
Background:	CD127 is a 60-90 kD type I transmembrane		
glycoprotein also known as IL-7 receptor α chain or			
	IL-7R α . It forms a heterodimer with the common γ		
	chain (γc or CD132) which is shared with the		
	receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and		
	IL-21. CD127 is expressed on immature B cells		
	through early pre-B stage, thymocytes (except		
	CD4/CD8 double positive thymocytes), peripheral T		
	cells, and bone marrow stromal cells. CD127 has		



	memory and effect with its receptor i mature and imma	be an useful marker for identifying etor T cells. The ligation of IL-7 s important for stimulation of ture T cells as well as immature B and development.
Form: Liquid		Isotype: Rat IgG2a, κ
Size:		Host: Rat
50Tests/100Tests/	/100Tests×2	
Reactivity: Mouse		Application: FCM
Concentration: 5	μL	Conjugation: AF488
SwissProt:	P16872	
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined

DEVELOP

for individual use.

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD127 Monoclonal Antibody (AF488 Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.