

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD127/IL-7RA Monoclonal
Antibody(PE Conjugated)

Catalog: DL20400F

Synonyms: Interleukin-7 receptor subunit alpha, IL-7R, IL-7
receptor subunit alpha, IL-7R subunit
alpha, IL-7R-alpha, IL-7RA, CD127

Background: CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7R α . It forms a heterodimer with the common γ chain (γ c or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has

been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

Form: Liquid

Isotype: Rat IgG2a, κ

Size: 25 μ g/100 μ g

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration: 0.2 mg/mL

Conjugation: PE

SwissProt: P16872

Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1

$\mu\text{g}/10^6\text{cells}$ in 100 μL volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD127 Monoclonal Antibody(PE Conjugated)[Used at $0.02 \mu\text{g}/10^6$ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}\text{C}$ and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.