

Manufacturer/S	upplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-80				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Anti-Mouse CD127/IL-7RA Monoclonal			
	Antibody(AF647 Conjugated)			
Catalog:	DL20408F			
Synonyms:	Interleukin-7 receptor subunit alpha,Il7r,IL-7			
	receptor subunit alpha,IL-7R subunit			
	alpha,IL-7R-alpha,IL-7RA,CD127			
Background:	CD127 is a 60-90 kD type I transmembrane			
glycoprotein also known as IL-7 receptor $\alpha$ chain or IL-7R $\alpha$ . It forms a heterodimer with the common $\gamma$				
				chain ( $\gamma c$ or CD132) which is shared with the
receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and				
	IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/CD8 double positive thymocytes), peripheral T			
cells, and bone marrow stromal cells. CD127 has				



	been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.		
Form: Liquid		Isotype: Rat IgG2a, κ	
Size: 25µg/100µg	5	Host: Rat	
Reactivity: Mouse		Application: FCM	
Concentration: 0.5 mg/mL		Conjugation: AF647	
SwissProt:	P16872		
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.		

## **Recommended Use:**


Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1

## **DEVELOP**

 $\mu g/10^6$  cells in 100  $\mu L$  volume].

## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD127 Monoclonal Antibody(AF647 Conjuaged)[Used at 0.2 µg/10<sup>6</sup> cells dilution]and Anti-Mouse CD3 Monoclonal Antibody(AF488 Conjuaged). Splenocytes stained with Anti-Mouse CD3 Monoclonal Antibody(AF488 Conjuaged)(right) are used as control.



## Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

www.dldevelop.com