

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-8	2732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	AF/LE Purified Anti-Mouse CD28 Monoclonal		
	Antibody		
Catalog:	DL20416F		
0			
Synonyms:	T-cell-specific surface glycoprotein		
	CD28,Cd28,CD28		
Background:	CD28 is a 44 kD glycoprotein, also known as Tp44		
	or T44. It is a member of the Ig superfamily,		
	expressed on thymocytes, most peripheral T cells, and NK cells. In association with CD80 (B7-1) and		
CD86 (B7-2), CD28 acts as the second signal for T			
	and NK cell activation and proliferation. The 37.51		
	antibody has been reported to augment in vitro T cell		
	proliferation and cytokine production, and promote		
	CTL development.		



Form: Liquid		Isotype: Syrian Hamster IgG
Size: 50µg/500µg/1mg		Host: Hamster
Reactivity: Mouse		<b>Application:</b> Activ, Block, FCM, IHC, IP
Concentration: 0.5 mg/mL		<b>Conjugation:</b> None (Purified antibody, Azide Free/Low endotoxin)
SwissProt:	P31041	
Buffer:	0.2 μm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or	

## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \ \mu g \ per \ 10^6 \ cells$  in 100  $\ \mu L$  volume or 100  $\ \mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

## **Storage:**

## **DEVELOP**.

Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.

