

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD28 Monoclonal Antibody(APC

Conjugated)

**Catalog:** DL20421F

**Synonyms:** T-cell-specific surface glycoprotein

CD28,Cd28,CD28

**Background:** CD28 is a 44 kD glycoprotein, also known as Tp44

or T44. It is a member of the Ig superfamily,

expressed on thymocytes, most peripheral T cells, and NK cells. In association with CD80 (B7-1) and CD86 (B7-2), CD28 acts as the second signal for T and NK cell activation and proliferation. The 37.51 antibody has been reported to augment in vitro T cell proliferation and cytokine production, and promote

CTL development.



Form: Liquid **Isotype:** Syrian Hamster IgG

Size: Host: Hamster

50Tests/100Tests/100Tests×2

**Reactivity:** Mouse **Application:** FCM

Concentration: 5 µL **Conjugation:** APC

SwissProt: P31041

Phosphate buffered solution, pH 7.2, containing **Buffer:** 

0.09% stabilizer and 1% protein protectant.


## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD28 Monoclonal Antibody(APC Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.





## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.