

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-8	FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD28 Monoclonal		
	Antibody(PE/Cyanine5.5 Conjugated)		
Catalog:	DL20424F		
Synonyms:	T-cell-specific surface glycoprotein		
	CD28,Cd28,CD28		
Daalamaundu	CD28 is a 44 kD glyconnotain also known as Tr44		
Dackground.	Background: CD28 is a 44 kD glycoprotein, also known as Tp44 or T44. It is a member of the Ig superfamily,		
	expressed on thymocytes, most peripheral T cells,		
and NK cells. In association with CD80 (B7-1) and $CD86$ (B7-2). $CD28$ gate as the second size of far T			
CD86 (B7-2), CD28 acts as the second signal for T and NK cell activation and proliferation. The 37.51			
	antibody has been reported to augment in vitro T cell		
	proliferation and cytokine production, and promote		
	CTL development.		



Form: Liquid		Isotype: Syrian Hamster IgG
Size: 20 Tests/50 Tests/100 Tests/200 Tests		Host: Hamster
Reactivity: Mouse		Application: FCM
Concentration:		Conjugation: PE/Cyanine5.5
SwissProt:	P31041	
Buffer:	PBS with 0.05% Proclin300, 1% BSA	

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.



This product is guaranteed up to one year from purchase.

