

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD40 Monoclonal Antibody(Biotin
Conjugated)

Catalog: DL20439F

Synonyms: Tumor necrosis factor receptor superfamily member
5, Cd40, B-cell surface antigen CD40, Bp50, CD40L
receptor, CD40, Tnfrsf5

Background: CD40 is a 48 kD type I transmembrane glycoprotein
also known as Bp50. It is a member of the tumor
necrosis factor receptor (TNFR) superfamily and is
expressed on B cells, basal epithelial cells,
macrophages, follicular dendritic cells, endothelial
cells, and a subset of CD34+ hematopoietic
progenitors. CD40 regulates B cell
development/maturation, Ig isotype switching and, in
combination with other signals such as IL-4, protects

B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.

Form: Liquid

Isotype: Rat IgG2a, κ

Size: 25 μ g/100 μ g

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration: 0.5 mg/mL

Conjugation: Biotin

SwissProt: P27512

Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.