

Manufaatuvar/Si	undier		
Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD40 Monoclonal		
	Antibody(PE/Cyanine5 Conjugated)		
Catalog:	DL20443F		
Synonyms:	Tumor necrosis factor receptor superfamily member 5,Cd40,B-cell surface antigen CD40,Bp50,CD40L receptor,CD40,Tnfrsf5		
Background:			
also known as Bp50. It is a member of the tumor necrosis factor receptor (TNFR) superfamily and is			
			expressed on B cells, basal epithelial cells,
macrophages, follicular dendritic cells, endothelial			
	cells, and a subset of CD34+ hematopoietic		
	progenitors. CD40 regulates B cell		
	development/maturation, Ig isotype switching and, in		
	combination with other signals such as IL-4, protects		
	comoniation with other signals such as 12-4, protects		



B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.

Form: Liquid		Isotype: Rat IgG2a, κ
Size: 50Tests/100Tests/100Tests×2		Host: Rat
Reactivity: Mouse		Application: FCM
Concentration: 5 µL		Conjugation: PE/Cyanine5
SwissProt:	P27512	
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined

DEVELOP

for individual use.

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD40 Monoclonal Antibody(PE/Cyanine5 Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.