

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD40 Monoclonal

Antibody(PerCP/Cyanine5.5 Conjugated)

**Catalog:** DL20446F

**Synonyms:** Tumor necrosis factor receptor superfamily member

5,Cd40,B-cell surface antigen CD40,Bp50,CD40L

receptor, CD40, Tnfrsf5

**Background:** CD40 is a 48 kD type I transmembrane glycoprotein

> also known as Bp50. It is a member of the tumor necrosis factor receptor (TNFR) superfamily and is

expressed on B cells, basal epithelial cells,

macrophages, follicular dendritic cells, endothelial

cells, and a subset of CD34+ hematopoietic

progenitors. CD40 regulates B cell

development/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects



B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.

Form: Liquid Isotype: Rat IgG2a, κ

Size: Host: Rat

50Tests/100Tests/100Tests×2

**Application:** FCM **Reactivity:** Mouse

Concentration: 5 µL Conjugation: PerCP/Cyanine5.5

SwissProt: P27512

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

## Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined



for individual use.

## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD40 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.