

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD1d Monoclonal Antibody(PE

Conjugated)

Catalog: DL20493F

Synonyms: Antigen-presenting glycoprotein

CD1d1,Cd1d1,CD1d.1,Cd1d1,Cd1.1

Background: CD1d is a type I transmembrane protein and member

of the MHC family, with a molecular weight ranging

from 43-49 kD, depending on the glycosylation degree. CD1d is expressed by antigen presenting cells such as dendritic cells, monocytes, macrophages

and B cells; also expressed by thymocytes and

intestinal epithelial cells. CD1d present glycolipids to iNKT cells that recognize them by their Vα14i TCR.

Form: Liquid Isotype: Rat IgG2b, κ



Size: 25µg/100µg Host: Rat

Reactivity: Mouse **Application:** FCM

Concentration: 0.2 mg/mL **Conjugation: PE**

SwissProt: P11609

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in 100 μL volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD1d Monoclonal Antibody(PE Conjugated)[Used at 0.2 μg/10⁶ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.





Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.