

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD210 Monoclonal Antibody(Biotin
Conjugated)

Catalog: DL20551F

Synonyms: Interleukin-10 receptor subunit alpha, IL10ra, IL-10
receptor subunit alpha,
IL-10R subunit
alpha, IL-10RA, CDw210a, Interleukin-10 receptor
subunit 1, IL-10R subunit 1, IL-10R1, CD210, IL10ra

Background: CD210 is a 90-110 kD IL-10 receptor. It is a class II
cytokine receptor expressed on thymocytes, T cells,
B cells, NK cells, monocytes and macrophages.
Ligand binding of CD210 induces Jak1 and Tyk,
resulting in STAT1 and STAT3 activation. IL-10
receptor stimulation results in the inhibition of
cytokine production and the costimulation of B cell

proliferation and differentiation. The only known ligand for this receptor is IL-10.

Form: Liquid

Isotype: Rat IgG1, κ

Size: 25 μ g/100 μ g

Host: Rat

Reactivity: Mouse

Application: FCM

Concentration: 0.5 mg/mL

Conjugation: Biotin

SwissProt: Q61727

Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:



Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.