

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD40 Monoclonal Antibody(Biotin

Conjugated)

Catalog: DL20557F

Synonyms: Tumor necrosis factor receptor superfamily member

5,Cd40,B-cell surface antigen CD40,Bp50,CD40L

receptor, CD40, Tnfrsf5

Background: CD40 is a 48 kD type I glycoprotein also known as

> BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells,

fibroblasts, and at low levels on plasma cells. CD40

has been reported to be involved in B cell differentiation, costimulation, isotype

class-switching, and protection of B cells from apoptosis. Additionally, CD40 is important for T



cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand). The 5C3 antibody has been reported to promote B cell proliferation in the presence of anti-IgM, IL-4 or PMA, partially blocking CD40 binding to CD40L, and B cells rescue from apoptosis.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: $25 \mu g/100 \mu g$ Host: Mouse

Reactivity: Human **Application:** FCM

Concentration: 0.5 mg/mL **Conjugation:** Biotin

SwissProt: P25942

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5 \mu g$ per 106 cells in 100 μl volume or 100 μl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.