

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD40 Monoclonal

Antibody(PerCP/Cyanine5.5 Conjugated)

Catalog: DL20565F

Synonyms: Tumor necrosis factor receptor superfamily member

5,Cd40,B-cell surface antigen CD40,Bp50,CD40L

receptor, CD40, Tnfrsf5

Background: CD40 is a 48 kD type I glycoprotein also known as

> BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages,

follicular dendritic cells, endothelial cells,

fibroblasts, and at low levels on plasma cells. CD40

has been reported to be involved in B cell differentiation, costimulation, isotype

class-switching, and protection of B cells from

apoptosis. Additionally, CD40 is important for T



cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand). The 5C3 antibody has been reported to promote B cell proliferation in the presence of anti-IgM, IL-4 or PMA, partially blocking CD40 binding to CD40L, and B cells rescue from apoptosis.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: Host: Mouse

20Tests/100Tests/100Tests×2

Application: FCM **Reactivity:** Human

Concentration: 5 µL Conjugation: PerCP/Cyanine5.5

SwissProt: P25942

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment.



Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD40 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.