

Manufacturer/Su	ıpplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Pro	vince, China			
TELE: 86-510-8	2732223 FAX: 86-510-82720101-8014			
Web : www.dldev	velop.com			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Anti-Human CD44 Monoclonal			
	Antibody(PE/Cyanine7 Conjugated)			
Catalog:	DL20575F			
Synonyms:	CD44 antigen,CD44,CDw44,Epican,Phagocytic			
	glycoprotein 1,PGP-1,Phagocytic glycoprotein			
	I,PGP-I,CD44,LHR, MDU2, MDU3, MIC4			
Background:	CD44 is a 80-95 kD glycoprotein also known as			
Dackground.	Hermes, Pgp1, H-CAM, or HUTCH. It is expressed			
on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44				
			expression increases from a low or mid level of	
		intensity to high expression levels. Thus, CD44 has		
			been reported to be a valuable marker for memory	
	cell subsets. CD44 is an adhesion molecule involved			



in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.

Form: Liquid		Isotype: Mouse IgG2a, κ	
Size:		Host: Mouse	
20Tests/100Te	ests/100Tests×2		
Reactivity: Human		Application: FCM	
Concentration: 5 µL		Conjugation: PE/Cyanine7	
SwissProt:	P16070		
Buffer:	-	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:



Human peripheral blood lymphocytes are stained with Anti-Human CD44 Monoclonal Antibody(PE/Cyanine7 Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.