

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD44 Monoclonal
Antibody(PE/Cyanine5.5 Conjugated)

Catalog: DL20576F

Synonyms: CD44 antigen, CD44, CDw44, Epican, Phagocytic
glycoprotein 1, PGP-1, Phagocytic glycoprotein
I, PGP-I, CD44, LHR, MDU2, MDU3, MIC4

Background: CD44 is a 80-95 kD glycoprotein also known as
Hermes, Pgp1, H-CAM, or HUTCH. It is expressed
on all leukocytes, endothelial cells, hepatocytes, and
mesenchymal cells. As B and T cells become
activated or progress to the memory stage, CD44
expression increases from a low or mid level of
intensity to high expression levels. Thus, CD44 has
been reported to be a valuable marker for memory
cell subsets. CD44 is an adhesion molecule involved

in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.

Form: Liquid

Isotype: Mouse IgG2a, κ

Size: 20 Tests/50 Tests/100 Tests/200 Tests

Host: Mouse

Reactivity: Human

Application: FCM

Concentration:

Conjugation: PE/Cyanine5.5

SwissProt: P16070

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ l staining volume or per 100 μ l of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.