

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-8	FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Human CD44 Monoclonal		
	Antibody(PerCP/Cyanine5.5 Conjugated)		
Catalog:	DL20577F		
Synonyms:	CD44 antigen,CD44,CDw44,Epican,Phagocytic glycoprotein 1,PGP-1,Phagocytic glycoprotein		
	I,PGP-I,CD44,LHR, MDU2, MDU3, MIC4		
Background:	Background: CD44 is a 80-95 kD glycoprotein also known as		
Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and			
		mesenchymal cells. As B and T cells become	
activated or progress to the memory stage, CD44			
expression increases from a low or mid level of			
	intensity to high expression levels. Thus, CD44 has		
	been reported to be a valuable marker for memory		
	cell subsets. CD44 is an adhesion molecule involved		



in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.

Form: Liquid		Isotype: Mouse IgG2a, κ	
Size:		Host: Mouse	
20Tests/100Te	sts/100Tests×2		
Reactivity: Human		Application: FCM	
Concentration: 5 µL		Conjugation: PerCP/Cyanine5.5	
SwissProt:	P16070		
Buffer:	-	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:



Human peripheral blood lymphocytes are stained with Anti-Human CD44 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.