

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Human CD54 Monoclonal Antibody(PerCP		
	Conjugated)		
Catalog:	DL20633F		
Synonyms:	Intercellular adhesion molecule		
	1,Icam1,MALA-2,MyD10,CD54,Icam-1		
Background:	CD54 is a 85-110 kD type I transmembrane protein		
	also known as ICAM-1. It is expressed on activated		
	endothelial cells, high endothelial venules, T and B		
	cells, monocytes/macrophages, granulocytes, and		
	dendritic cells. The expression of ICAM-1 can be		
	released from the cell surface. CD54 plays a role in		
	cellular adhesion and is involved in inflammation		
	and leukocyte extravasation. CD54 has also been		
	shown to be the major cellular receptor for		
	rhinovirus. ICAM-1 binds to CD11a/CD18 (LFA-1),		



κ

CD11b/CD18 (Mac-1), CD11c/CD18 (p150, 95) as well as hyaluronan and fibrinogen.

Form: Liquid		Isotype: Mouse IgG1,
Size: 20 Tests/50 Tests/100 Tests/200 Tests		Host: Mouse
Reactivity: Human		Application: FCM
Concentration:		Conjugation: PerCP
SwissProt:	P05362	
Buffer:	PBS with 0.05% Proclin300, 1% BSA	

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ l staining volume or per 100 μ l of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

DEVELOP.

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

