

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD20 Monoclonal Antibody(APC

Conjugated)

**Catalog:** DL20644F

**Synonyms:** B-lymphocyte antigen CD20,MS4A1,B-lymphocyte

> surface antigen B1, Bp35, Leukocyte surface antigen Leu-16, Membrane-spanning 4-domains subfamily A

member 1,MS4A1,CD20

**Background:** CD20 is a 33-37 kD, four transmembrane spanning

> protein, also known as B1 and Bp35. CD20 is expressed on pre-B-cells, resting and activated B cells (not plasma cells), some follicular dendritic cells, and at low levels on a T cell subset. CD20 is heavily phosphorylated on activated B cells and malignant B cells. Homo-oligomeric complexes of CD20 are thought to form Ca2+ conductive ion



channels in the plasma membrane of B cells. The CD20 molecule is involved in B-cell activation and is associated with various Src family kinases (Lyn, Lck, Fyn). It exists in a complex with MHC class I and II, CD53, CD81, and CD82.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: Host: Mouse

20Tests/100Tests/100Tests×2

**Reactivity:** Human **Application:** FCM

Concentration: 5 µL **Conjugation:** APC

SwissProt: P11836

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

## Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined



for individual use.

## Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD20 Monoclonal Antibody(APC Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.