

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Purified Anti-Human CD29 Monoclonal Antibody Product name:

DL20676F **Catalog:** 

**Synonyms:** Integrin beta-1,ITB1,Itgb1,Fibronectin receptor

subunit beta, VLA-4 subunit beta, CD29

**Background:** CD29 is a 130 kD single chain type I glycoprotein

> also known as integrin β1, VLA-β chain, or gpIIa. It is broadly expressed on a majority of hematopoietic and non-hematopoietic cells, including leukocytes (although at low level on granulocytes), platelets, fibroblasts, endothelial cells, epithelial cells, and mast cells. CD29 is a member of the integrin family. It is non-covalently associated with integrin  $\alpha 1-\alpha 6$

chains to form VLA-1 to VLA-6 molecules,

respectively. Integrins, which include CD29, bind to several cell surface (e.g. VCAM-1, MadCAM-1) and



extracellular matrix molecules. CD29 acts as a fibronectin receptor and is involved in a variety of cell-cell and cell-matrix interactions.

Form: Liquid **Isotype:** Mouse IgG1, κ

**Size:** 25µg/100µg **Host:** Mouse

**Reactivity:** Human **Application:** FCM,IHC,IF,IP

**Concentration:** 0.5 mg/mL Conjugation: Unconjugated

**SwissProt:** P05556

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer.

## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.5 \mu g$  per  $10^6$  cells in  $100 \mu L$  volume or  $100 \mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



## **Storage:**

Keep as concentrated solution.

Store at  $2\sim8$ °C. Do not freeze.

This product is guaranteed up to one year from purchase.