

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD31 Monoclonal

Antibody(PerCP/Cyanine5.5 Conjugated)

Catalog: DL20697F

Synonyms: Platelet endothelial cell adhesion

molecule, PECAM1, PECAM-1, EndoCAM, GPIIA', P

ECA1,CD31

Background: CD31 is a 130-140 kD type I transmembrane

> glycoprotein also known as platelet endothelial cell adhesion molecule-1 (PECAM-1) or Endocam. It is expressed on monocytes, platelets, granulocytes, endothelial cells and lymphocyte subsets. CD31 has been reported to bind CD38 and be involved in

> wound healing, angiogenesis, and cellular migration

in an inflammatory situation.



Form: Liquid **Isotype:** Mouse IgG1, κ

Size: **Host:** Mouse

20Tests/100Tests/100Tests×2

Reactivity: Human **Application:** FCM

Concentration: 5 µL Conjugation: PerCP/Cyanine5.5

SwissProt: P16284

Phosphate buffered solution, pH 7.2, containing **Buffer:**

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use

Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD31 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.





Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.