

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD147 Monoclonal

Antibody(PE/Cyanine7 Conjugated)

Catalog: DL20743F

Synonyms: Basigin, BSG, 5F7, Collagenase stimulatory

> factor, Extracellular matrix metalloproteinase inducer, EMMPRIN, Leukocyte activation antigen M6,OK blood group antigen, Tumor cell-derived collagenase stimulatory factor, TCSF, CD147

Background: CD147, also known as neurothelin or basigin, is a

> member of the Ig superfamily. It is a 55-65 kD type I transmembrane glycoprotein which is primarily expressed on leukocytes, erythrocytes, platelets, and

endothelial cells. CD147 is reported to have a

function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain



endothelia.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: Host: Mouse

20Tests/100Tests/100Tests×2

Reactivity: Human **Application:** FCM

Concentration: 5 µL Conjugation: PE/Cyanine7

SwissProt: P35613

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:



Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.