

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD38 Monoclonal

Antibody(PE/Cyanine5.5 Conjugated)

Catalog: DL20768F

Synonyms: ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase

1,CD38,2'-phospho-ADP-ribosyl

cyclase,2'-phospho-cyclic-ADP-ribose

transferase, ADP-ribosyl cyclase 1, ADPRC 1, Cyclic

ADP-ribose hydrolase 1,cADPr hydrolase

1,T10,CD38

Background: CD38 is a 45 kD type II transmembrane glycoprotein

> also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is

expressed at high levels on plasma cells and activated



T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: Host: Mouse

20Tests/100Tests/100Tests×2

Reactivity: Human **Application:** FCM

Concentration: 5 µL Conjugation: PE/Cyanine5.5

SwissProt: P28907

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.