

Manufacturer/S	upplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-8	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Purified Anti-Human CD47 Monoclonal Antibody			
Catalog:	DL20772F			

Synonyms: Leukocyte surface antigen CD47,Cd47,Integrin-associated protein,IAP

Background: CD47 also known as Rh-associated protein, gp42, integrin-associated protein (IAP), and neurophilin, is a 42-52 kD member of the immunoglobulin superfamily containing a five-pass transmembrane attachment. Two splice variants have been described in the cytoplasmic tail, the shorter form is expressed in bone-marrow-derived cells, endothelial cells, and fibroblasts while the longer form is expressed by neural tissues. CD47 expression is widely distributed in hematopoietic cells including thymocytes, T cells, B cells, monocytes, platelets, and erythrocytes as



well as epithelial cells, endothelial cells, fibroblasts, and neural tissues. CD47 functions as an adhesion molecule and thrombospondin receptor and is non-covalently associated with  $\beta$ 3 integrins CD51/CD61, CD41/CD61. Thrombospondin is a ligand for CD47; in the absence of CD47 mice show defects in host defense and  $\beta$ 3 integrin-dependent ligand binding, migration, and cellular activation. CD47 is also part of the Rh complex on erythrocytes.

Form: Liquid		<b>Isotype:</b> Mouse IgG1, κ	
Size: 25µg/100µg		Host: Mouse	
Reactivity: Human		Application: FCM	
Concentration: 0.5 mg/mL		Conjugation: Unconjugated	
SwissProt:	Q08722		
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.		

## **Recommended Use:**

Each lot of this antibody is quality control tested by flow

www.dldevelop.com



cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.5 \ \mu g \ per \ 10^6 \ cells$  in 100  $\mu L$  volume or 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

## **Storage:**

Keep as concentrated solution. Store at 2~8°C. Do not freeze. This product is guaranteed up to one year from purchase.

