

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD48 Monoclonal

Antibody(PE/Cyanine5.5 Conjugated)

Catalog: DL20792F

Synonyms: CD48 antigen, Cd48, BCM1 surface

antigen, BLAST-1, HM48-1, MRC OX-45 surface

antigen, SLAMF2, sgp-60, CD48

Background: CD48 is a 40-47 kD GPI-anchored membrane

> protein, also known as Blast-1 and HuLy-m3. It is a member of the CD2 family that contains 2 IgSF domains and is widely expressed on both resting and activated hematopoietic cells with the exception of granulocytes, platelets, and erythrocytes. CD48 binds to CD2 at a considerably (>100-fold) lower affinity than CD58. It is thought to contribute to T cell

activation. The cytoplasmic tail of CD48 has been



shown to bind to the kinases Lck and Fyn.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: 20 Tests/50 Tests/100 Host: Mouse

Tests/200 Tests

Reactivity: Human **Application:** FCM

Concentration: Conjugation: PE/Cyanine5.5

SwissProt: P09326

PBS with 0.05% Proclin300, 1% BSA Buffer:

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µl staining volume or per 100 µl of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

Keep as concentrated solution.



Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.