

Manufacturer/Supplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Pro	vince, China	
TELE: 86-510-82732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>		
Email: info@dldevelop.com.cn service@dldevelop.com.cn		
Product name:	Anti-Human CD55 Monoclonal Antibody(AF488	
	Conjugated)	
Catalog:	DL20825F	
_		
Synonyms:	ms: Complement decay-accelerating factor,CD55,CR,	
	DAF	
Background:	CD55 is a 60-70 kD glycosylphosphatidylinositol	
_	(GPI)-anchored single chain glycoprotein also known	
	as decay-accelerating factor (DAF). It is expressed	
	on hematopoietic cells including erythrocytes and	
	many non-hematopoietic cells. CD55 accelerates the	
	dissociation of the components of the C3-convertases	
	(namely C2a from C4b in the C4bC2a complex, a	
	C3-convertase of the classical pathway, and factor	
	Bb from the C3bBb complex, a C3-convertase of the	
	alternative pathway) to protect cells from	
	1 37 1	



inappropriate damage caused by autologous complement. CD55 has been reported to reduce the efficiency of NK cell lysis and induce signal transduction in T cells. CD55 has also been shown to interact with CD97 and bind to Coxsackie and Echovirus.

Form: Liquid	Isotype: Mouse IgG1, κ
Size: 20 Tests/50 Tests/100	Host: Mouse
Tests/200 Tests	
Reactivity: Human	Application: FCM
Concentration:	Conjugation: AF488
SwissProt: P08174	

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ l staining volume or per 100 μ l of whole blood). Please check your vial before the experiment. Since applications vary, the



appropriate dilutions must be determined for individual use.

Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD55 Monoclonal Antibody (AF488 Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

